

Reply to Weiss: Tree-ring stable oxygen isotopes suggest an increase in Asian monsoon rainfall at 4.2 ka BP

Bao Yang^{a,b,1}, Chun Qin^a, Achim Bräuning^c, Timothy J. Osborn^d, Valerie Trouet^e, Fredrik Charpentier Ljungqvist^{f,g,h} Jan Esperⁱ , Lea Schneider , Jussi Grießinger , Ulf Büntgen , Sergio Rossi , Guanghui Dong , Mi Yan , Liang Ning , Jianglin Wang^a, Xiaofeng Wang^a®, Beixi Fan^a, Suming Wang^r, Jürg Luterbacher^s, Edward R. Cook^t, Nils Chr. Stenseth^{u, 1}®

Weiss (1) raises interesting points on our article (2). He observes that the Delingha tree-ring δ^{18} O record matches KM-A speleothem δ^{18} O record from the Mawmluh Cave in India, that defines the 4.2-ka event's global-type stratum (3). This event is also manifested as a multicentennial drought in the Iranian Gol-E-Zard speleothem record (4). Weiss further argues that the Delingha record is also consistent with the 4.2-ka anomaly recorded in lowresolution proxy records from northern China despite inherent limitations of these proxies.

Our Delingha $\delta^{18}\text{O}$ record shows a persistent transition following gradual drying that, instead, occurred during 4 ka to 3.5 ka. This is associated with generally wet conditions ca. 4.5 ka to 4.0 ka (Fig. 1), in contrast with the KM-A δ^{18} O record. We thus do not observe such a close correspondence between the two records in Weiss's (1) figure 1A and want to exercise caution in drawing conclusions based on a limited number of proxy records with considerable age uncertainties.

To complement the four speleothem δ^{18} O records presented by Weiss (1), we assembled a total of 24 additional records from eastern Asia and India covering Holocene hydroclimate dynamics. In western China, only one out of three speleothem records from Kesang cave, KS06-A-H (5), has a good dating accuracy to resolve centennial drought anomalies and shows dry climate conditions at 4.2 ka (see ref. 6). In northern China, all records except Dongshiya and Zhenzhu indicate an anomalously wet interval around 4.2 ka (6). Both exceptions are characterized by few and uncertain dating points and a coarse sampling resolution (6). In southern China, investigating the full set of records yields no clear picture (6). However, a composite based on six high-precision (Fig. 1) records shows generally wet conditions at 4.2 ka.

In India, two more-recent replicated speleothem δ¹⁸O records from Mawmluh Cave with exceptional chronologic constraints and high sampling resolution have challenged the interpretation of KM-A (6). One of these records, ML.1, shows wet conditions at 4.1 ka to 4.0 ka (7), and the subsequent long-term multicentennial drying is consistent with our Delingha δ^{18} O record and with a long-term drying trend in a recent highresolution speleothem δ^{18} O record from Sahiya Cave in northern India (8).

In conclusion, our well-replicated Delingha δ^{18} O record does not support a significant hydroclimate transition in our study region around 4.2 ka, or the notion that this rapid climate deterioration should be regarded as generalized climatic transition from the Middle to

Late Holocene. There is the need for further development of accurate, replicated, and high-resolution proxy data, based on dense sampling, a mechanistic understanding of climatic controls, and representative chronological control points to elucidate the spatial-temporal dynamics of Holocene climate and cultural responses. Our Delingha δ^{18} O record makes considerable progress toward those aims.

ACKNOWLEDGMENTS. We thank Liangcheng Tan, Yanjun Cai, Xunlin Yang, and Gayatri Kathayat for providing part of the speleothem data used in this study. B.Y. is funded by the National Natural Science Foundation of China (Grants 42130511 and 41888101).

Author affiliations: a Key Laboratory of Desert and Desertification, Northwest Institute of Eco-Environment and Resources, Chinese Academy of Sciences, Lanzhou 730000, China; bChinese Academy of Sciences Center for Excellence in Tibetan Plateau Earth Sciences, Chinese Academy of Sciences, Beijing 100101, China; Institute of Geography, Friedrich-Alexander-University Erlangen-Nürnberg, 91058 Erlangen, Germany; dClimatic Research Unit, School of Environmental Sciences, University of East Anglia, Norwich NR4 7TJ, United Kingdom; ^eLaboratory of Tree-Ring Research, University of Arizona, Tucson, AZ 85721; fDepartment of History, Stockholm University, 106 91 Stockholm, Sweden; gBolin Centre for Climate Research, Stockholm University, 106 91 Stockholm, Sweden; hSwedish Collegium for Advanced Study, 752 38 Uppsala, Sweden; ¹Department of Geography, Johannes Gutenberg University, 55099 Mainz, Germany; ¹Department of Geography, Justus-Liebig-University, D-35390 Giessen, Germany; ^kDepartment of Geography, University of Cambridge, CB2 3EN Cambridge, United Kingdom; Dendrosciences Group, Swiss Federal Research Institute for Forest, Snow, and Landscape Research, CH-8903 Birmensdorf, Switzerland; "Czech Globe Global Change Research Institute, Czech Academy of Sciences, 60300 Brno, Czech Republic; "Department of Geography, Faculty of Science, Masaryk University, 61137 Brno, Czech Republic; ^oDépartement des Sciences Fondamentales, Université du Québec à Chicoutimi, Chicoutimi, QC G7H 2B1, Canada; PCollege of Earth and Environmental Sciences, Lanzhou University, Lanzhou 730000, ^qKey Laboratory of Virtual Geographic Environment of Ministry of Education, School of Geography Science, Nanjing Normal University, Nanjing 210023, China; State Key Laboratory of Lake Science and Environment, Nanjing Institute of Geography and Limnology, Chinese Academy of Sciences, Nanjing 210008, China; Science and Innovation Department, World Meteorological Organization, CH-1211 Geneva, Switzerland; ^tTree Ring Laboratory, Lamont Doherty Earth Observatory of Columbia University, Palisades, NY 10964; and ^uCentre for Ecological and Evolutionary Synthesis, Department of Biosciences, University of Oslo, N-0316 Oslo, Norway

Author contributions: B.Y. designed research; B.Y., C.Q., A.B., T.J.O., V.T., F.C.L., J.E., L.S., J.G., U.B., S.R., G.D., M.Y., L.N., J.W., X.W., B.F., S.W., J.L., E.R.C., and N.C.S. performed research; and B.Y. wrote the paper in close interactions with all authors.

The authors declare no competing interest.

Copyright © 2022 the Author(s). Published by PNAS. This article is distributed under Creative Commons Attribution-NonCommercial-NoDerivatives License 4.0 (CC BY-NC-ND).

¹To whom correspondence may be addressed: Emails: yangbao@lzb.ac.cn or n.c. stenseth@mn.uio.no.

Published May 13, 2022.




Fig. 1. The Delingha tree-ring δ^{18} O record (red line) and normalized speleothem δ^{18} O records collected from different parts of continental Asia spanning 5 ka to 3 ka. To place the 4.2-ka event in high-resolution and high-precision context, we constrained our analyses to speleothem records with coverage of 5 ka to 3 ka. A regional composite for southern China is shown, which is derived by averaging the six speleothem 618O series from Dongge Cave, Xianglong Cave, Wuya Cave, Sanbao Cave 43, Heshang Cave, and Shennong Cave using the Z-score method. Each of the six speleothem records has a temporal resolution better than 20 y, at least five U-Th ages, and dating precision higher than the 60-y average age error (2a) in the 5- to 3-ka interval. Prior to averaging, each of the six records is first linearly interpolated annually, and then their long-term linear trends in the common period 5 ka to 3 ka are removed to highlight climate fluctuations on multidecadal to centennial timescales before normalization. See ref. 6 for details about each stalagmite record employed in the calculation. The Liu-li Cave in northern China and Kesang Cave (KS06-A-H) in western China speleothem δ^{18} O records are intentionally included for comparison in that both records have a dating point around 4.2 ka (6). The 100-point low-pass filters are shown for Kesang Cave (KS06-A-H), Liu-li Cave, and southern China records. The Shennong Cave and Mawmluh Cave (ML.1) speleothem s^{18} O records have biannual and subannual resolution, and, therefore, we plot their unsmoothed series for comparison. All horizontal lines represent the long-term average calculated over the common period 5 ka to 3 ka. The vertical dashed blue line indicates the Middle to Late Holocene transition as defined by the Delingha δ^{18} O record via trend-point analysis (2). The gray bar covers the period 3.97 ka to 4.26 ka that is demonstrated as the 4.2-ka event in the midlatitudes of the Northern Hemisphere by the Iranian Gol-E-Zard speleothem record (4).

- H. Weiss, The East Asian summer monsoon, the India summer monsoon, and the midlatitude westerlies at 4.2 ka BP. *Proc. Natl. Acad. Sci. U.S.A.*, 10.1073/pnas.2200796119 (2022).
 B. Yang *et al.*, Long-term decrease in Asian monsoon rainfall and abrupt climate change events over the past 6,700 years. *Proc. Natl. Acad. Sci. U.S.A.* 118, e2102007118 (2021).
 M. Walker *et al.*, Subdividing the Holocene Series/Epoch: Formalization of stages/ages and subseries/subepochs, and designation of GSSPs and auxiliary stratotypes. *J. Quat. Sci.* 34, 173-186 (2019).
 S. A. Carolin *et al.*, Precise timing of abrupt increase in dust activity in the Middle East coincident with 4.2 ka social change. *Proc. Natl. Acad. Sci. U.S.A.* 116, 67-72 (2019).
 H. Cheng *et al.*, The climatic cyclicity in semiarid-arid central Asia over the past 500,000 years. *Geophys. Res. Lett.* 39, L01705 (2012).
 B. Yang *et al.*, SI Appendix for Reply to Weiss. Zenodo. https://doi.org/10.5281/zenodo.6473441.

- G. Kathayat et al., Evaluating the timing and structure of the 4.2 ka event in the Indian summer monsoon domain from an annually resolved speleothem record from Northeast India. Clim. Past 14, 1869-1879 (2018).
- 8. G. Kathayat et al., The Indian monsoon variability and civilization changes in the Indian subcontinent. Sci. Adv. 3, e1701296 (2017).