

<https://doi.org/10.1038/s44168-025-00327-1>

Arctic driftwood proposal for durable carbon removal

Ulf Büntgen, Clive Oppenheimer, Mirek Trnka, Michael Kempf, Ian Holman, Tito Arosio, Tatiana Bebchuk & Jan Esper

Check for updates

Various geoengineering approaches have been proposed for carbon dioxide (CO₂) removal but their viability at scale remains unclear. Here, we consider the natural behaviour of driftwood, the warming-induced acceleration of sea-ice loss and tree growth, as well as the stability of cellulose in subfossil wood under cold-anoxic conditions, to introduce the concept of sinking timber from the boreal forest for durable CO₂ sequestration at the deep Arctic Ocean floor.

Capture and storage of atmospheric CO₂ offer a means to stabilise climate alongside emission-reduction efforts. However, it is estimated that over 10 gigatonnes (Gt) of CO₂ would have to be removed and sequestered each year over the 21st century to mitigate legacy effects of anthropogenic greenhouse gas emissions and compensate for those sources expected to remain hard to decarbonise^{1–3}. While reductions of fossil fuel burning must be prioritised at national and international levels^{3,4}, different hybrid nature-engineering technologies have been recommended to capture and store CO₂ from the Earth's atmosphere. Although frequently presented as viable strategies for mitigating the effects of greenhouse gas emissions^{1,2}, many approaches face questions regarding their scalability and the risk of counterproductive consequences for humans and the environment⁵.

Among proposed solutions is 'Wood Vaulting' (WV) or 'Wood Harvesting and Storage' (WHS)^{1,2,5}, which involves burying woody biomass in engineered enclosures that inhibit decomposition under anaerobic or frozen conditions, thereby ideally sequestering carbon on multi-millennial or even longer timescales. A prototype Wood Vault Unit (WVU) of 1 ha spatial extent and 20 m soil depth could store around 10⁵ m³ of timber, which is equivalent to approximately 0.1 Mt CO₂. It would therefore take annual construction of 10⁴ WVU to operate at 1 Gt yr⁻¹ of CO₂ removal, corresponding to a roughly 25% increase in global logging (currently around 4 × 10⁹ m³ of wood annually⁶). Substantial ecological and societal trade-offs can be expected from operating at such scale, including lasting impacts on soil carbon and mycorrhizal networks, biodiversity loss, and co-emissions associated with deforestation, transportation and vault construction^{5,7}. Further, the putative benefits of WV would be offset if only a fraction of methane generated from decaying wood reaches the atmosphere^{8,9}.

Here, we examine the natural occurrence and behaviour of driftwood from the boreal forest to introduce a variant of WHS that would involve durable carbon storage on the deep, near anoxic floor of the Arctic Ocean.

Driftwood solution for carbon sequestration

The circumpolar boreal forest zone stretches across northern North America and Eurasia, from Alaska and northern Canada through Scandinavia and across the Siberian taiga. Characterised by cold climates, slow growing conifers, widespread peatlands, extensive permafrost soils, and gigantic river systems^{10–12}, the world's largest terrestrial biome also represents an enormous carbon pool^{13,14}, with as much as 10³ Gt (10¹⁸ g) of carbon stored in living trees, dead wood, soils and peat¹⁵. Unlike wildland tropical forests, the estimated carbon stocks of boreal forest ecosystems are likely to increase under global warming¹⁶, though whether the taiga as a whole becomes a net source or sink of carbon under warming remains unclear¹⁵. Parts of the boreal forest export large quantities of organic matter to riparian zones and fluvial networks, which ultimately reach the Arctic Ocean via surface runoff, riverbank erosion and mass wasting^{17,18}. This drainage includes substantial but unquantified amounts of coarse woody material, known as driftwood¹⁹, some of which accumulates in the vast delta systems of large boreal rivers and along Arctic coastlines^{20,21}.

Riverbank erosion strongly controls the amount of natural driftwood transported to the Arctic Ocean (Fig. 1A–C). In open ocean conditions, intact stems typically remain buoyant for 1 yr depending on species, but when entrained in sea ice they can be transported for several years before being released¹⁹. Timescales for wood to sink to the deep floor of the Arctic Ocean depend on density contrast, ocean depth and currents but are likely significantly shorter than floating time. This natural process is accelerating due to the combined effects of warming-induced permafrost thaw and forest expansion^{22,23}, as well as rapid sea-ice loss and increased river discharge^{24,25} (Fig. 1D, E).

A few very large river systems drain broad sectors of the circumpolar boreal forest and deliver the majority of terrestrial freshwater and dissolved and particulate organic carbon to the Arctic Ocean^{17,18,25,26}. The most important catchments (and their discharge) of the Russian taiga from west to east are those of the Ob (400 km³ yr⁻¹), Yenisey (590 km³ yr⁻¹), Lena (540 km³ yr⁻¹), and Kolyma (70 km³ yr⁻¹), while the Mackenzie in Canada (290 km³ yr⁻¹) and Yukon in Alaska (210 km³ yr⁻¹) drain most of the taiga in northern North America^{17,26}. Collectively they represent an estimated 11% of global freshwater runoff²⁶. While increasing their discharge rates, and hence the amount of driftwood that can reach the Arctic Ocean, recent anthropogenic warming is also affecting the capacity of individual trees and entire forest ecosystems to sequester CO₂ from the atmosphere. Total ring width and maximum latewood density values of conifers across the boreal forests of northern North America and Eurasia have been increasing since a period of reduced growth at the end of the 20th century^{27,28} (Fig. 1F).

A key aspect of our thought experiment on using driftwood to sequester atmospheric carbon is the negligible rate of wood decay after sinking (Fig. 1C). Extremely low decay rates under near anoxic and freezing conditions^{29,30} suggest the deep Arctic Ocean floor would be highly suited for




Fig. 1 | Arctic amplification and driftwood solution. **A** natural erosion and tree tipping, as well as **(B)** driftwood accumulation along the Indigirka river in north-eastern Siberia (taken by Ulf Büntgen in July 2016). **(C)** Underwater logs on the ocean floor of the northwest continental shelf of the Chinese Sea⁵¹. **D** changes in sea-ice concentration >60 and >66° North (red and orange, respectively) expressed as total annual sums of sea-ice cover anomalies⁵², **(E)** changes in vapour pressure (warmer and wetter conditions) >60 and >66° North (dark and light blue,

respectively) expressed as average hPa anomalies⁵³, and **(F)** changes in tree-ring width (TRW; light green) and maximum latewood density (MXD; dark green) expressed as mean and median (thin dashed lines) timeseries after ‘signal-free age-dependent spline’ detrending of eight undisturbed boreal forest sites in northern North America and northern Eurasia (<https://climexp.knmi.nl> and <https://www.monostar.org>).

long-term storage. This is supported by measurements of circa 200 living and relict, dry-dead and subfossil trees from different cold, oxic and anoxic environments in the European Alps (e.g., talus, lakes and peat), which revealed no systematic decline in α -cellulose content over the past 8000 years³¹ (Fig. 2). The centennial to multi-millennial scale stability of wood carbon is further corroborated by decades of worldwide dendrochronological research on dry-dead and subfossil wood samples from historical buildings, archaeological excavations, and sediments spanning Holocene and even Pleistocene contexts³². While the composition and deposition of boreal driftwood should be confirmed, we expect the combination of low temperature, reduced oxygen and limited wood-borer activity to characterise large parts of the Arctic shelf and deep basin^{33–35}.

Conclusion and projection

Though widely discussed (and frequently criticised)^{36–38}, planting trees for carbon removal and storage has limited impact beyond their lifespan (captured by the adage “grow fast and die young”)³⁹. Evidence also suggests that afforestation of Arctic tundra is likely to result in net warming due to reduced surface albedo³⁸, negating perceived climate change mitigation benefits of high-latitude tree planting on previously unforested terrain. Instead, we suggest further exploration of the potential of harvesting and rafting large quantities of boreal timber into the Arctic Ocean for CO₂ removal and multi-millennial scale storage (Fig. 3). Given access to carbon rich, and economically unimportant boreal conifer trees with short transit

routes to large river systems, combined with efficient monocultural reforestation practices, the cold Arctic Ocean could store vast quantities of carbon from Siberia and northern North America where biodiversity is low and the risk of wildfires high⁴⁰. The burning-induced succession of boreal forests has almost tripled during the first two decades of the 21st century as the biome became warmer⁴¹.

To achieve significant CO₂ drawdown, we propose, for the purposes of our thought experiment, three units of circa 10,000 km² (comparable to the size of Lake Onega in northwestern Russia near the Finnish border) for extensive harvesting and reforestation along each of the five main rivers and their tributaries in Russia, Alaska and Canada: Ob, Yenisey, Lena, Yukon, and Mackenzie. Due to high fire risk (and low human population), these regions carry ~10–30 t/ha of larch, pine or spruce timber for harvesting (at decreasing mass per unit area with increasing latitude). Taking 15 t/ha stand carbon content, annual logging and rafting of circa 180,000 km² timber could remove up to 1 Gt/y of CO₂. The total area of harvesting would represent around 1% of the boreal forest zone, comparable with the area consumed annually by wildfires^{42,43}. All target regions should be even-aged, biodiversity-poor and fire-prone monocultural coniferous stands of low economic and cultural value. If logging is mainly carried out in winter, access may be facilitated by extensive ice roads, clearing can be performed on solid ground, and timber can be placed directly on the frozen rivers. Mulching small branches and other wooden remains can decrease fire risk, increase soil development, and enhance nutrient availability.




Fig. 2 | Wood preservation and carbon sequestration. **A** Alpha-cellulose content in 17 living and circa 183 relict, dry-dead and subfossil larch (*Larix decidua* Mill.) and pine (*Pinus cembra* L.) trees from the Austrian and Swiss Alps between 1950 and 2400 m asl, where wood preservation is promoted by near freezing conditions³¹. Brown horizontal bars show the timespan of the individual wood samples after precise cross-dating (x-axis) and the median α -cellulose content expressed in

percentage and calculated from five-year blocks. The dashed line is the mean and suggests that there are no long-term effects of possible wood decay on α -cellulose content in living, dry-dead and subfossil trees over the past 8000 years (6980 BCE to 2015 CE). **B** Box plots summarise data for each millennium over much of the Holocene. We also measured 26.4% (± 7.16) of remaining α -cellulose in a radiocarbon-dead subfossil tree trunk from northern Greenland (not shown).

Natural and silvicultural reforestation is likely to sequester most CO₂ during the first few decades of forest regeneration^{44,45}. Such a multi-year, seasonal cycle of harvesting, sinking and replanting will always capture more CO₂ than any form of natural taiga succession in which trees grow slower and will either burn or decompose afterwards. Potential removal rates, however, can be expected to vary substantially between biogeographic zones, and boreal forests are less productive (but more durable) than those in warmer climates^{44,45}. It should be further noted that the boreal rivers and their vast delta systems^{19–21}, together with large parts of the circumpolar coastlines of northern North America and Eurasia already contain significant amounts of driftwood⁴⁶.

Although our thought experiment should not distract from the priority of reducing greenhouse gas emissions, with continued economic growth undermining efforts to meet the Paris Agreement targets, carbon removal proposals are increasingly relevant⁴⁷. As with other means for carbon capture and removal, our silvicultural proposal is not without caveats and requires further interdisciplinary scientific investigation. We recognise significant issues must be evaluated carefully in developing and refining our concept not least concerning land ownership by indigenous peoples, infrastructure and market value, topography, hydrology, accessibility, biodiversity, and productivity of different harvest units in the boreal forest zone, as well as the species-specific sinking potential of driftwood under changing sea-ice conditions, and the locations of its final deposition in more or less anoxic parts of the Arctic Ocean floor. Undesirable environmental impacts that might arise include the release of phenols and other wood chemicals

during both controlled and uncontrolled river rafting, and ocean sinking, while large quantities of floating timber may threaten riverine and maritime traffic. Geo-political questions concerning different cost factors and ownership rights of the Arctic Ocean floor would also need to be addressed, including whether seabed driftwood storage should be accounted as part of the terrestrial or marine environment, with implications for carbon sink and source budgeting at national and international scales (and hence carbon credit incentivisation). Rigorous cost-benefit modelling with a comprehensive agent-perspective for environmental and societal impact assessments is also needed (Fig. 3). Such a model must accurately address multi-scalar, cross-cultural and cross-functional/sectoral⁴⁸ tensions between the norm and value-based institutions of indigenous forest user groups and the market cost and revenue generation processes of the logging and climate mitigation industry^{49,50}. A refined model is expected to define ecologically, economically and politically suitable harvesting practices, logging terrains and shipping routes (Fig. 3).

While logging at a desirable scale could hypothetically be achieved by Russia alone, we imagine a coordinated circumpolar effort that complements other mitigation strategies. Following scientific and indigenous guidance, the incentive for Moscow, Ottawa and Washington to start considering a viable concept of using driftwood to sequester atmospheric carbon could be twofold: Reductions of greenhouse gas emissions to mitigate the effects of anthropogenic climate and environmental change, in tandem with fiscal profit from carbon credit points, and international reputation for sustainable nature-based geoeengineering.




Fig. 3 | Driftwood carbon storage model with agent-perspective. **A** Circumpolar boreal forest zone with large river systems, and the extent of burnt boreal forest between 2002 and 2020 that amounts to circa 1,835,00 km² (red areas)^{42,43}. **B** Least-cost analysis of a boreal forest patch with suitable timber harvesting parameters and optimal driftwood transportation conditions along the closest river to the nearest ocean²⁴. Floating time is calculated as average downstream river run-off velocity and depending on rafting style and wood amount. An ecological buffer zone has been included around the nearest administrative centre from which labour and logistics are directed. The simplified model design includes an agent-perspective³⁵, in which

the ability for the exogenous (e.g., harvesting for wood products and wood vaulting, and maintenance for carbon offsetting) and endogenous (e.g., cultural, herding, etc) demand for forest services to be met by spatial production depends on factors such as forest productivity, land ownership, infrastructure, human resources and the decisions of modelled agents, informed by their values, objectives and perceptions of future monetary and non-monetary value of land. **C** Pictures of natural driftwood erosion, tree tipping and driftwood rafting, as well as Indigenous people at the Indigirka river in northeastern Siberia (all taken by Ulf Büntgen in July 2016).

Data availability

No datasets were generated or analysed during the current study.

Ulf Büntgen^{1,2,3} ✉, Clive Oppenheimer¹, Mirek Trnka^{2,4},
Michael Kempf^{1,5}, Ian Holman^{2,6}, Tito Arosio^{1,7}, Tatiana Bebchuk¹ &
Jan Esper^{2,8}

¹Department of Geography, University of Cambridge, Cambridge, UK.

²Global Change Research Institute (CzechGlobe), Czech Academy of Sciences, Brno, Czech Republic. ³Department of Geography, Faculty of Science, Masaryk University, Brno, Czech Republic. ⁴Department of Agrosystems and Bioclimatology, Faculty of Agronomy, Mendel University, Brno, Czech Republic. ⁵Quaternary Geology, Department of Environmental Sciences, University of Basel, Basel, Switzerland.

⁶Cranfield University, Bedfordshire, UK. ⁷Forest Dynamics, Swiss Federal Institute for Forest, Snow and Landscape Research WSL, Birmensdorf, Switzerland. ⁸Department of Geography, Johannes Gutenberg University, Mainz, Germany.

✉ e-mail: ulf.buentgen@geog.cam.ac.uk

Received: 2 October 2025; Accepted: 18 December 2025;

Published online: 03 January 2026

References

- Zeng, N. et al. Carbon sequestration via wood harvest and storage: an assessment of its harvest potential. *Clim. Change* **118**, 245–257 (2013).
- Zeng, N. et al. 3775-year-old wood burial supports “wood vaulting” as a durable carbon removal method. *Science* **385**, 1454 (2024).
- IPCC. *Climate change 2023: Synthesis Report Contribution of Working Groups I, II and III to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change* (eds Core Writing Team, Lee, H. & Romero, J.) (IPCC, 2023).
- Smith, S. M. et al. *The State of Carbon Dioxide Removal 2nd edn.* <https://doi.org/10.17605/OSF.IO/F85QJ> (University of Manchester, 2024).
- Zeng, N. & Hausmann, H. Wood Vault: remove atmospheric CO₂ with trees, store wood for carbon sequestration for now and as biomass, bioenergy and carbon reserve for the future. *Carbon Balance Manag.* **17**, 2 (2022).
- FAO. *Food and Agricultural Organization of the United Nations.* <https://www.fao.org/faostat/en> (FAO, 2025).
- Peng, L., Searchinger, T. D., Zions, J. & Waite, R. The carbon costs of global wood harvests. *Nature* **620**, 110–115 (2023).
- Covey, K. R. & Megonigal, J. P. Methane production and emissions in trees and forests. *N. Phytol.* **222**, 35–51 (2019).
- Barba, J. et al. Methane emissions from tree stems: a new frontier in the global carbon cycle. *N. Phytol.* **222**, 18–28 (2019).
- Bonan, G. B. & Shugart, H. H. Environmental factors and ecological processes in boreal forests. *Annu. Rev. Ecol. Syst.* **20**, 1–28 (1989).
- Esseen, P.-A., Ehnström, B., Ericson, L. & Sjöberg, K. Boreal Forests. *Ecol. Bull.* **46**, 16–47 (1997).
- McClelland, J. W., Déry, S. J., Peterson, B. J., Holmes, R. M. & Wood, E. F. A pan-arctic evaluation of changes in river discharge during the latter half of the 20th century. *Geophys. Res. Lett.* **33**, L06715 (2006).
- Yu, Z. C. Northern peatland carbon stocks and dynamics: a review. *Biogeosciences* **9**, 4071–4085 (2012).
- Pan, Y. et al. A large and persistent carbon sink in the world’s forests. *Science* **333**, 988–993 (2011).
- Bradshaw, C. J. & Warkentin, I. G. Global estimates of boreal forest carbon stocks and flux. *Glob. Planet. Change* **128**, 24–30 (2015).
- Hogan, J. A. et al. Anthropomes and forest carbon responses to global change. *Plants People Planet* **7**, 1027–1042 (2025).
- Holmes, R. M. et al. A. Seasonal and annual fluxes of nutrients and organic matter from large rivers to the Arctic Ocean and surrounding seas. *Estuaries Coasts* **35**, 369–382 (2012).
- McClelland, J. W. et al. Particulate organic carbon and nitrogen export from major Arctic rivers. *Glob. Biogeochem. Cycles* **30**, 629–643 (2016).
- Hellmann, L. et al. Dendro-provenancing of Arctic driftwood. *Quat. Sci. Rev.* **162**, 1–11 (2017).
- Stadie, C. et al. Large driftwood accumulations along arctic coastlines and rivers. *Sci. Rep.* **15**, 32500 (2025).
- Büntgen, U., Kirdyanov, A. V., Hellmann, L., Nikolayev, A. & Tegel, W. Cruising an archive: on the palaeoclimatic value of the Lena Delta. *Holocene* **24**, 627–630 (2014).
- Friggens, N. L. et al. Positive rhizosphere priming accelerates carbon release from permafrost soils. *Nat. Commun.* **16**, 3576 (2025).
- Dial, R. J. et al. Arctic sea ice retreat fuels boreal forest advance. *Science* **383**, 877–884 (2024).
- Kolar, T. et al. Predicted sea-ice loss likely terminates Iceland’s driftwood supply by 2060 CE. *Glob. Planet. Change* **213**, 103834 (2022).
- Shiklomanov, A. et al. River freshwater flux to the Arctic Ocean. in *Arctic Hydrology, Permafrost and Ecosystems* (Springer, 2020).
- Holmes, R. M., Shiklomanov, A. I., Tank, S. E., McClelland, J. W. & Tretiakov, M. River Discharge. in *Arctic Report Card 2015.* <http://www.arctic.noaa.gov/Report-Card> (2015).
- Briffa, K. R. et al. Reduced sensitivity of recent tree-growth to temperature at high northern latitudes. *Nature* **391**, 678–682 (1998).
- Büntgen, U., Kirdyanov, A. V., Krusic, P. J., Shishov, V. V. & Esper, J. Arctic aerosols and the ‘Divergence Problem’ in dendroclimatology. *Dendrochronologia* **67**, 125837 (2021).
- Russell, M. B. et al. Quantifying carbon stores and decomposition in dead wood: a review. *For. Ecol. Manag.* **350**, 107–128 (2015).
- Röllig, R. et al. Wood decay under anoxia by the brown-rot fungus *Fomitopsis pinicola*. *Nat. Commun.* **16**, 7352 (2025).
- Arosio, T. et al. No long-term decay in α-cellulose of living and relict trees from the European Alps over the past 8000 years. *Paleoceanogr. Paleoclimatol.* **40**, e2025PA005170 (2025).
- Reinig, F. et al. Precise date for the Laacher See eruption synchronizes the Younger Dryas. *Nature* **595**, 66–69 (2021).
- Hellmann, L. et al. Tracing the origin of Arctic driftwood. *J. Geophys. Res.-B* **118**, 68–76 (2013).
- Borges, L. M. S., Merkelbach, L. M. & Cragg, S. M. Biogeography of wood-boring crustaceans (Isopoda: Limnoriidae) established in European coastal waters. *PLoS ONE* **9**, e109593 (2014).
- Björdal, C. G. & Dayton, P. K. First evidence of microbial wood degradation in the coastal waters of the Antarctic. *Sci. Rep.* **10**, 12774 (2020).
- Bastin, J.-F. et al. The global tree restoration potential. *Science* **365**, 76–79 (2019).
- Moyano, J. et al. Unintended consequences of planting native and non-native trees in treeless ecosystems to mitigate climate change. *J. Ecol.* **112**, 2480–2491 (2024).
- Kristensen, J. Å. et al. Tree planting is no climate solution at northern high latitudes. *Nat. Geosci.* **17**, 1087–1092 (2024).
- Büntgen, U. et al. Limited capacity of tree growth to mitigate the global greenhouse effect under predicted warming. *Nat. Commun.* **10**, 2171 (2019).
- Kirdyanov, A. V. et al. Long-term ecological consequences of forest fires in the continuous permafrost zone of Siberia. *Environ. Res. Lett.* **15**, 034061 (2020).
- Chandanpurkar, H. A. et al. Unprecedented continental drying, shrinking freshwater availability, and increasing land contributions to sea level rise. *Sci. Adv.* **11**, eadx0298 (2025).
- Scholten, R. C., Veraverbeke, S., Chen, Y. & Randerson, J. T. Spatial variability in Arctic-boreal fir regimes influenced by environmental and human factors. *Nat. Geosci.* **17**, 866–873 (2024).
- Loboda, T. V. et al. *Arctic Boreal Annual Burned Area, Circumpolar Boreal Forest and Tundra, V2, 2002–2022 (Version 2).* <https://doi.org/10.3334/ORNLDAA/2328> (2024).
- Bernal, B., Murray, L. T. & Pearson, T. R. H. Global carbon dioxide removal rates from forest landscape restoration activities. *Carbon Balance Manag.* **13**, 22 (2018).
- Robinson, N. et al. Protect young secondary forests for optimum carbon removal. *Nat. Clim. Change* **15**, 793–800 (2025).
- Sendrowski, A., Wohl, E., Hilton, R., Kramer, N. & Ascough, P. Wood-based carbon storage in the Mackenzie River Delta: the world’s largest mapped riverine wood deposit. *Geophys. Res. Lett.* **50**, e2022GL100913 (2023).
- Jiang, J., Shi, S. & Raftery, A. E. Mitigation efforts to reduce carbon dioxide emissions and meet the Paris Agreement have been offset by economic growth. *Commun. Earth Environ.* **6**, 823 (2025).
- Rahman, H. M. T. & Pigford, A. Institutionalism in sustainable environmental management: an increasing relevance for institutional diversity. in *Institutional Diversity and Environmental Sustainability* (eds Rahman, H.M.T. & Pigford, A.) (CRC Press, 2025).
- Rahman, H. M. T. & Natcher, D. Addressing gaps in integrative water-energy-food-forest (WEFF) nexus governance. *Environ. Sci. Policy* **172**, 104195 (2025).
- Booth, A. L. & Skelton, N. W. There’s a conflict right there: Integrating indigenous community values into commercial forestry in the Tl’azt’en First Nation. *Soc. Nat. Resour.* **24**, 368–383 (2011).
- Chu, S. et al. Degradation condition and microbial analysis of waterlogged archaeological wood from the second shipwreck site on the northwestern continental slope of the South China Sea. *npj Herit. Sci.* **13**, 10 (2025).
- Titchner, H. A. & Rayner, N. A. The Met Office Hadley Centre sea ice and sea surface temperature data set, version 2: 1. Sea ice concentrations. *J. Geophys. Res. Atmos.* **119**, 2864–2889 (2014).
- Harris, I., Osborn, T. J., Jones, P. & Lister, D. Version 4 of the CRU TS monthly high-resolution gridded multivariate climate dataset. *Sci. Data* **7**, 109 (2020).
- Kempf, M. et al. Carbon dioxide removal: code and simulation. *Zenodo.* <https://doi.org/10.5281/zenodo.17112515> (2025).
- Blanco, V. et al. The effect of forest owner decision-making, climatic change and societal demands on land-use change and ecosystem service provision in Sweden. *Ecosyst. Serv.* **23**, 174–208 (2017).

Acknowledgements

This study was supported by the AdAgriF project: “Advanced methods of greenhouse gases emission reduction and sequestration in agriculture and forest landscape for climate change mitigation” (CZ.02.01.01/00/22_008/0004635), the ERC Advanced Grant (882727; Monostar), and the ERC Synergy Grant (101118880; Synergy-Plague). We are thankful to colleagues in Brno, Cambridge and Mainz for stimulating discussions.

Author contributions

U.B. and J.E. initiated and conceived the study. U.B. wrote the manuscript together with C.O., M.T., I.H. and J.E., whereas M.K. was responsible for the model aspect of the study. T.A. provided cellulose data and T.B. contributed to discussion and revision.

Competing interests

The authors declare no competing interests.

Additional information

Correspondence and requests for materials should be addressed to Ulf Büntgen.

Reprints and permissions information is available at <http://www.nature.com/reprints>

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2026